

Codimension one spheres which are null homotopic.

Laurence R. Taylor

Grove and Halperin [3] introduced a notion of *taut immersions*. Terng and Thorbergsson [5] give a slightly different definition and showed that taut immersions are a simultaneous generalization of taut immersions of manifolds into Euclidean spaces or spheres, and some interesting embeddings constructed by Bott and Samelson [1]. They go on to prove many theorems about such immersions. One particularly intriguing result, Theorem 6.25, concerned codimension one, null homotopic, tautly embedded spheres. Using a result of Ruberman, [4], they proved in many cases that this sphere had to be a distance sphere, that is, the image of a standard sphere under the exponential map, a generalization of a theorem of Chern and Lashof [2]. Here we observe that the methods of Terng–Thorbergsson and Ruberman suffice to classify tautly immersed, null homotopic, codimension one spheres. Informally one may say that the examples produced by Terng and Thorbergsson in [5] are all that there are. Precisely, we have

Theorem 1. *Let N^n be a complete Riemannian manifold and let $\phi: S \rightarrow N$ be a tautly immersed, codimension one sphere which is null homotopic. Then there exists a taut point $q \in N$ and a real number l , $0 < l < r_{conj}(q)$, such that S is the image of the sphere of radius l in $T_q N$ under the exponential map. Here $r_{conj}(q)$ denotes the conjugate radius at the point $q \in N$.*

Remark. If we fix $q \in N$ and a real number l , $0 < l < r_{conj}(q)$, then the image of the sphere of radius l under the exponential map from $T_q N$ is a taut immersion if and only if q is taut. This is Theorem 6.23, p. 219, of Terng and Thorbergsson [5]. It follows from the definition of conjugate radius that $exp_q: T_q N \rightarrow N$ is an immersion when restricted to the ball of radius l and from Theorem 6.23 [5] it follows that the spheres parallel to the boundary are all taut.

To explain the result of Ruberman eluded to above, we introduce some notation. A locally flat, codimension one sphere in M which disconnects M displays M as a connected sum, $M = M_1 \# M_2$. We call this sphere a *connected sum sphere*. If we assume a codimension one, locally flat sphere is null homotopic, it must be a connected sum sphere. One obvious way to get null homotopic, connected sum spheres is for at least one of M_1 or M_2 to be a homotopy sphere. It is natural to wonder if there are any others.

Ruberman [4] gives examples of null homotopic connected sum spheres for which neither M_1 nor M_2 is a homotopy sphere. He also gives a nearly necessary and sufficient criterion to have a null homotopic, connected sum sphere, [4,Thm. 1]. We observe that a more vigorous use of Ruberman’s main lemma provides a complete characterization. This

Partially supported by the N.S.F.

characterization shows that Ruberman's list of examples is complete. The characterization is given by the following theorem.

Theorem 2. *A connected sum sphere in M , a connected, paracompact, Hausdorff, m -dimensional topological manifold, with or without boundary, is null homotopic if and only if either (1) or (2) below holds, where $M = M_1 \# M_2$.*

- (1) *At least one of M_1 or M_2 is a homotopy m -sphere.*
- (2) *Neither M_1 nor M_2 is a homotopy sphere, but there exist relatively prime integers k_1 and k_2 , both greater than 1, so that each M_i is a simply connected $\mathbf{Z}[\frac{1}{k_i}]$ -homology m -sphere.*

Remarks. We repeat Ruberman's remark that case (2) can not hold if $m \leq 4$, but there are infinitely many examples of (2) in each dimension ≥ 5 . We emphasize that case (1) must occur unless M is a compact, simply connected manifold without boundary which is a rational homology m -sphere with torsion in its homology containing at least two distinct primes. We also remark that Ruberman already proved Theorem 2 in the case M is simply connected and this is the only case we use in the proof of Theorem 1.

The author would like to thank D. Ruberman and G. Thorbergsson for useful conversations.

1. The proof of Theorem 2.

To state Ruberman's main lemma requires further notation and conventions. To form the connected sum requires embeddings $\iota_i: B^m \rightarrow M_i$. Then $M_1 \# M_2$ is formed by removing the interiors of $\iota_i(B^m)$ and identifying the boundaries. If $M_1 \vee M_2$ denotes the wedge M_1 and M_2 then there is a natural map $M_1 \# M_2 \rightarrow M_1 \vee M_2$ obtained by pinching $i(S^{m-1})$ to a point and identifying $M_i/\iota_i(B^m)$ with M_i via an inverse to the map $M_i \rightarrow M_i/\iota_i(B^m)$ which is the identity outside a small neighborhood of B^m . Let $\hat{B}_i \subset M_i$ be a larger ball containing $\iota_i(B^m)$ so that $\hat{B}_i - \iota_i(B_i) = S^{m-1} \times [0, 1]$. There are maps $M_i \rightarrow S^m$ which pinch all of $M_i - \hat{B}_i$ to a point. Composition yields a map $M_1 \# M_2 \rightarrow M_1 \vee M_2 \rightarrow S^m \vee S^m$. Orient $S^{m-1} = \partial B^m$ and use the inward normal last of ∂B^m to orient the balls in M_1 and M_2 . This orients the two S^m as well.

Now suppose the connected sum sphere is null homotopic. This means that i extends to a map of the disk $I: D^m \rightarrow M_1 \# M_2$. The map I yields two maps $I_i: S^m \rightarrow M_i$. Define the degree of the map I_i as the degree of the composite $S^m \xrightarrow{I_i} M_i \rightarrow S^m$ where we orient the first S^m from D^m using the inward normal last from the orientation on S^{m-1} . Ruberman's main lemma then becomes

Lemma 3. (Ruberman [4, Lemma 3]). *With notation and conventions as above,*

$$\deg I_1 + \deg I_2 = 1 .$$

For completeness we give a proof. The following diagram commutes

$$\begin{array}{ccccc} M_1 \# M_2 & \rightarrow & M_1 \vee M_2 & \rightarrow & M_i \\ \downarrow & & \downarrow & & \downarrow \\ S^m \# S^m & \rightarrow & S^m \vee S^m & \rightarrow & S^m \end{array}$$

It follows that it suffices to prove the result for the case $M_1 = M_2 = S^m$. Orient $S^m \# S^m$ so that the map $\mathbf{Z} \simeq H_m(S^m \# S^m) \rightarrow H_m(S^m \vee S^m) \simeq \mathbf{Z} \oplus \mathbf{Z}$ sends 1 to $(1, -1)$. Given two null homotopies, $I, J: D^m \rightarrow S^m \# S^m$ of the connected sum sphere, they differ by an element in $\pi_m(S^m \# S^m)$ and the map $\mathbf{Z} \simeq \pi_m(S^m \# S^m) \rightarrow H_m(S^m \vee S^m; \mathbf{Z}) \simeq \mathbf{Z} \oplus \mathbf{Z}$ sends 1 to $(1, -1)$. Hence the sum of the degrees is constant. The evident null homotopy which collapses the connected sum sphere in $S^m \# S^m$ using the first S^m has degrees $(1, 0)$, so that difference is 1. ■

Two further lemmas will be required.

Lemma 4. *If X is a simply connected, $\mathbf{Z}[\frac{1}{k}]$ -acyclic space, then $\pi_{m-1}(X)$ is k -torsion, $m \geq 2$.*

As Ruberman observes, this is an elementary consequence of Serre's mod $\mathcal{C}$ theory. He also gives a proof of this next lemma as his Lemma 2.

Lemma 5. *If $f: S^m \rightarrow M^m$, $m > 1$, has degree k , $k \neq 0$, then $\pi_1(M)$ is finite of order dividing k . It further follows that M is a $\mathbf{Z}[\frac{1}{k}]$ -homology m -sphere.*

Remark 6. Note that a simply connected homology sphere is a homotopy sphere. If $k = \pm 1$ in Lemma 5, M is a homotopy sphere even if $m = 1$. If M is not closed, compact and orientable, then automatically $k = 0$.

First we prove that the connected sum sphere in case (1) or in case (2) is null homotopic. The proof presented is essentially Ruberman's. In case (1), the connected sum sphere is null homotopic because $M_i - D^m$ is a homotopy disk if M_i is a homotopy sphere. In case (2), recall that $m \geq 5$. Note that $M_i - D^m$ is $\mathbf{Z}[\frac{1}{k_i}]$ -acyclic and the map $S^{m-1} \xrightarrow{i} M_1 \# M_2$ factors through each $M_i - D^m$. By Lemma 4, the class in $\pi_{m-1}(M_1 \# M_2)$ represented by the connected sum sphere is both k_1 torsion and k_2 torsion. Since k_1 and k_2 are relatively prime, i is null homotopic.

Now suppose the connected sum sphere is null homotopic and that neither side is a homotopy sphere. We must show case (2) holds.

Step 1. We have $m > 1$ and there exist relatively prime numbers k_1 and k_2 , both greater than 1 such that each M_i is a $\mathbf{Z}[\frac{1}{k_i}]$ -homology m -sphere. The fundamental group of M_i is finite of order dividing k_i .

Proof: From Lemma 5, Remark 6 and our assumption that neither side is a homotopy sphere, it follows that neither $\deg I_i$ can be ± 1 . It follows from Lemma 3 that neither $\deg I_i$ can be 0 either. It follows from Remark 6 that M_i is closed, compact, orientable. If $m = 1$, $M_i = S^1$ which is a contradiction. Hence $m > 1$. Let k_i be the absolute value of $\deg I_i$ and note that each k_i is greater than 1. It follows from Lemma 3 that the k_i are relatively prime. It follows from Lemma 5 that each M_i has finite fundamental group of order dividing k_i and is a $\mathbf{Z}[\frac{1}{k_i}]$ -homology m -sphere. ■

If M is a rational homology sphere with finite fundamental group, let $r(M)$ denote the order of $\pi_1(M)$ and let $\ell(M)$ denote the order of the direct sum of the torsion subgroups of the homology. Step 1 can be summarized as

Summary. If the connected sum sphere in $M^m = M_1 \# M_2$ is null homotopic with M connected and neither M_i a homotopy sphere, then each M_i is a rational homology m -sphere, $(\ell(M_1), \ell(M_2)) = 1$, and $(r(M_1), r(M_2)) = 1$. Since neither M_i is a homotopy sphere, $m > 1$ and $\ell(M_i) \cdot r(M_i) > 1$.

Consider the following construction. Let $\widetilde{M}_i$ denote the universal cover of M_i and let Y_i be the connected sum of $\widetilde{M}_i$ and $(r(M_i) - 1)$ copies of M_{3-i} .

Step 2. Y_i is not a homotopy sphere.

Proof: Since M_{3-i} is not a homotopy sphere, neither is Y_i if $r(M_i) > 1$. But if $r(M_i) = 1$, $Y_i = M_i$ is not a homotopy sphere either. ■

The next two formulae follow from the Mayer–Vietoris and the van–Kampen theorems.

$$\begin{aligned} (5) \quad & \ell(Y_i) = \ell(\widetilde{M}_i) \cdot \ell(M_{3-i})^{r(M_i)-1} \\ (6) \quad & \pi_1(Y_i) = \pi_1(M_{3-i}) * \cdots * \pi_1(M_{3-i}) \quad r(M_i) - 1 \text{ times.} \end{aligned}$$

Step 3. The connected sum sphere in $Y_i \# M_{3-i}$ is null homotopic.

Proof: The manifold $Y_i \# M_{3-i}$ is the total space of a cover of $M_i \# M_{3-i}$: $\pi: Y_i \# M_{3-i} \rightarrow M_i \# M_{3-i}$ denotes the covering map. Check that π of the connected sum sphere in $Y_i \# M_{3-i}$ is the connected sum sphere in $M_i \# M_{3-i}$. Since $m \geq 2$, the induced map $\pi_*: \pi_{m-1}(Y_i \# M_{3-i}) \rightarrow \pi_{m-1}(M_i \# M_{3-i})$ is an injection. ■

The Final Step: $\pi_1(M_i) = 0$: i.e. $r(M_i) = 1$.

Proof: Suppose $r(M_i) > 1$. By Steps 2 and 3, the Summary applies to $Y_i \# M_{3-i}$. Hence $\ell(Y_i)$ and $\ell(M_{3-i})$ are relatively prime, so $\ell(M_{3-i}) = 1$ by (5). It follows from the Summary then that $r(M_{3-i}) > 1$. Since $\pi_1(Y_i)$ is finite, (6) implies $r(M_i) = 2$ and hence $r(Y_i) = r(M_{3-i})$. This is a contradiction since $(r(Y_i), r(M_{3-i})) = 1$. Hence $r(M_i) = 1$. ■

2. The proof of Theorem 1.

We begin with three results we need in the proof. First observe that more mileage is available from the proof of Theorem 2.5, p. 188 of [5].

Theorem 7. Let $\phi: M \rightarrow N$ be a taut immersion and fix a point $x \in M$. Assume M is connected. Let $\pi: \tilde{N} \rightarrow N$ be the cover for which there is a choice of $y \in \tilde{N}$ such that the subgroup $\pi_*(\pi_1(\tilde{N}, y)) \subset \pi_1(N, \phi(x))$ equals the subgroup $\phi_*(\pi_1(M, x)) \subset \pi_1(N, \phi(x))$. Let $\tilde{\phi}: M \rightarrow \tilde{N}$ be the unique lift with $\tilde{\phi}(x) = y$. Then $\tilde{\phi}$ is an embedding.

Proof: Note that $P(\tilde{N}, \tilde{\phi} \times q)$ is connected since M is and $\tilde{\phi}$ induces an isomorphism on π_1 . Next note that $P(\tilde{N}, \tilde{\phi} \times q)$ is a component of $P(N, \phi \times \pi(q))$. If we choose $q \in \tilde{N}$ as Terng and Thorbergsson do on page 188, then the energy function is a perfect Morse function on $P(\tilde{N}, \tilde{\phi} \times q)$. The result follows just as in [5]. ■

Proposition 8. *Let $\phi: M \rightarrow N$ be a map, let $\pi: \tilde{N} \rightarrow N$ be a cover, and let $\tilde{\phi}: M \rightarrow \tilde{N}$ be a map covering ϕ . Then ϕ is a taut immersion if and only if $\tilde{\phi}$ is.*

Proof: It follows from the structure of covering spaces that ϕ is an immersion if and only if $\tilde{\phi}$ is. Next note that an immersion $\psi: X \rightarrow Y$ is taut if and only if the energy function on the path space $P(Y, \psi \times q)$ is a perfect Morse–Bott function for all points $q \in Y$. One direction is Theorem 2.9, p. 193, in [5]. The other direction follows from the remark on page 183 of [5] that the energy function on the path space is Morse at all non-focal points, plus the observation that a proper Morse–Bott function which is Morse is a proper Morse function.

Now the path space for ϕ is the disjoint union of various path spaces

$$P(N, \phi \times p) = \coprod_{p_\alpha \in \pi^{-1}(p)} P(\tilde{N}, \tilde{\phi} \times p_\alpha) .$$

Since the energy function on each $P(\tilde{N}, \tilde{\phi} \times p_\alpha)$ is just the restriction of the energy function on $P(N, \phi \times p)$, the result is immediate. ■

Remark. G. Thorbergsson has shown the author an elementary proof of Proposition 8 that does not need Theorem 2.9.

Finally we need a generalization of Theorem 6.25 [5].

Proposition 9. *Let $\psi: S \rightarrow M$ be a taut embedding and assume $\psi(S)$ bounds a simply connected, $\mathbf{Z}/2\mathbf{Z}$ homology ball $B \subset M$. Then B is a ball and there exists a taut point $p \in B \subset M$ and a real number $l > 0$ such that $\psi(S)$ is the image of the sphere of radius l in $T_p M$ and l is less than $r_{conj}(p)$.*

Proof: We indicate the changes needed from the proof of Theorem 6.25 on pages 219 and 220 of [5]. Following Terng and Thorbergsson, pick $p \in B \subset M$ so that there is a geodesic perpendicular to $\psi(S)$ with first focal point p at a distance l from $\psi(S)$. We further assume l is minimal with this property. Observe that the space $P(B, \psi \times p)$ fits into a fibration sequence

$$\Omega B \rightarrow P(B, \psi \times p) \xrightarrow{\pi} S .$$

Since B is simply connected and mod 2 acyclic, it follows from the Serre spectral sequence that ΩB is also mod 2 acyclic. It then follows from the Serre spectral sequence that π_* induces an isomorphism in mod 2 homology. Hence $P(B, \psi \times p)$ has no homology below dimension $n-1$. Argue just as in Terng and Thorbergsson that the multiplicity of the focal point p is $n-1$, so all the geodesics starting perpendicularly to $\psi(S)$ meet at a distance l in the point p . Furthermore, all of these geodesics minimize distance between p and $\psi(S)$. Hence the conjugate radius at p is bigger than l . Furthermore no two of these geodesics intersect in B since they minimize distance, so B is the ball of radius l centered at p . It follows from [5, Thm. 6.23, p. 219] that p is taut. ■

To fix notation for the proof of Theorem 1, let $\pi: \tilde{N} \rightarrow N$ be the universal cover. Since $\phi: S^{n-1} = S \rightarrow N$ is null homotopic, ϕ lifts to the universal cover via $\tilde{\phi}: S \rightarrow \tilde{N}$. By Theorem 7, $\tilde{\phi}$ is an embedding and it remains null homotopic. Hence by Theorem 2,

there is a simply connected, $\mathbf{Z}/2\mathbf{Z}$ homology ball, $B \subset \tilde{N}$, with boundary $\partial B = \tilde{\phi}(S)$. The embedding $\tilde{\phi}$ is taut by Proposition 8. Applying Proposition 9, we now have a taut point $\tilde{q} \in \tilde{N}$ and a real number l such that $\tilde{\phi}(S)$ is the distance sphere of radius l centered at $\tilde{q}$ with $l < r_{conj}(\tilde{q}) = r_{conj}(q)$, where q denotes the image of $\tilde{q}$ in N . It follows that $\phi(S)$ is the image of the sphere of radius l in $T_q N$ under the exponential map. Finally apply Proposition 8 again: since $\tilde{q} \in \tilde{N}$ is taut, q is taut in N as required by Theorem 1.

Remark. The hypothesis in Theorem 1 that the immersion ϕ is null homotopic is sometimes unnecessary. By [5,Thm. 2.5], if $\pi_1(N) = 0$ then ϕ is an embedding and in many cases, the $\mathbf{Z}/2\mathbf{Z}$ cohomology ring will force one of the pieces in the connected sum decomposition to be a simply connected $\mathbf{Z}/2\mathbf{Z}$ homology ball. Then Proposition 9 will show that ϕ is null homotopic. Examples include all simply connected manifolds with the $\mathbf{Z}/2\mathbf{Z}$ cohomology of the rank 1 symmetric spaces, $H^*(S^n; \mathbf{Z}/2\mathbf{Z})$, $H^*(CP^n; \mathbf{Z}/2\mathbf{Z})$, $H^*(HP^n; \mathbf{Z}/2\mathbf{Z})$ and the $\mathbf{Z}/2\mathbf{Z}$ cohomology of the Cayley projective plane. Products of such spaces are also examples.

Remark. One can generalize Ruberman's theorem to show that if the order of $\phi \in \pi_{n-1}(N)$ is odd, then the embedding $\tilde{\phi}$ bounds a simply connected $\mathbf{Z}/2\mathbf{Z}$ homology ball, so again Proposition 9 will show that ϕ is null homotopic.

Bibliography

- [1] R. Bott and H. Samelson, *Applications of the theory of Morse to symmetric spaces*, Amer. J. Math. **80** (1958), 964–1029, Corrections in **83** (1961) 207–208.
- [2] S.-S. Chern and R. K. Lashof, *On the total curvature of immersed manifolds*, Amer. J. Math. **79** (1957), 306–318.
- [3] K. Grove and S. Halperin, *Elliptic isometries, condition (C) and proper maps*, Arch. Math. (Basel) **56** (1991), 288–299.
- [4] D. Ruberman, *Null homotopic spheres of codimension one*, “Tight and Taut Submanifolds” (T. E. Cecil and S.-S. Chern, eds.), MSRI Publications, vol. 32, Cambridge Univ. Press, Cambridge, 1997, pp. 229–232.
- [5] C.-L. Terng and G. Thorbergsson, *Taut immersions into complete Riemannian manifolds*, “Tight and Taut Submanifolds” (T. E. Cecil and S.-S. Chern, eds.), MSRI Publications, vol. 32, Cambridge Univ. Press, Cambridge, 1997, pp. 181–228.

Department of Mathematics, University of Notre Dame, Notre Dame, IN 46556

email address: *taylor.2@nd.edu*

URL: *<http://www.nd.edu/~taylor>*